

Forest Health Protection



Numbered Report 08-06

May 2008

2007 NORTH IDAHO DOUGLAS-FIR TUSSOCK MOTH TRAPPING SYSTEM REPORT

**Lee Pederson, Entomologist,
Doug Wulff, Biological Science Technician
USDA Forest Service, Northern Region Coeur d'Alene**

**Jeffrey Fidgen, Lands Program Manager,
Brian Burkhead, GIS Specialist,
Tom Eckburg, Lands Program Specialist,
Idaho Department of Lands**

INTRODUCTION

Idaho's Douglas-fir tussock moth (DFTM) Early Warning System (EWS) uses a series of pheromone trap sites to identify increasing populations prior to undesirable tree defoliation, a system modified after Daterman et al. (1979). The Idaho Department of Lands (IDL) maintains a network of trap sites from Coeur d'Alene south to Moscow and east to Harvard. Region 1 of the US Forest Service (USFS-R1) maintains sites from Potlatch to Lucille. A portion of these sites is monitored annually.

At each site, five pheromone-baited sticky traps are installed to monitor the flight of male moths. An average trap catch of 25 moths is the threshold used to indicate where heavy defoliation may occur the following year. Follow up sampling is then conducted in these areas to pinpoint injurious population densities (Daterman et al. 1979) and to apply treatments, if necessary.

METHODS

2007 Trapping Results

A total of 151 sites were monitored in northern Idaho during 2007. The mean trap capture was 0.43 moths per trap, up from 0.39 and 0.06 moths per trap in 2006 and 2005 respectively. None of the sites had trap captures exceeding 25 moths per trap. Two trap sites monitored by IDL had mean captures exceeding 5 moths per trap. Mason Butte (plot ID 800) and Little Plummer Creek (plot ID 803), both within the Coeur d'Alene Indian Reservation, had trap catches of 7.25 and 5.80 moths per trap respectively (Figure 1, Appendix 1). None of the traps monitored by USFS exceeded 5 moths per trap, although one site, Bargamin Creek (plot ID 5-4), had a mean trap capture of 4.6 moths per trap (Figure 2, Appendix 2).



Defoliation

The most recent outbreak in north Idaho began in 2000, resulting in three years of defoliation on state and private land between Plummer and Moscow, and in adjacent Clearwater National Forest lands (Fig. 4). Outbreaks of DFTM have typically occurred in this general area about every 8-10 years since the 1940's. Prior to the 2000 outbreak, visible defoliation occurred for one year during 1986. Both outbreaks were preceded by increasing numbers of trap captures (Randall 2002) (Fig. 3). This year's aerial detection survey showed very little DFTM-caused defoliation.

Larval Surveys

Larval surveys were performed at 45 of the 120 plots trapped by IDL in 2007 (Appendix 1), and all plots had low populations. Larval population surveys were conducted using the sequential sampling methods described by Mason (1978).

Conclusions

The DFTM, EWS has been effective at predicting outbreaks in northern Idaho. The two latest outbreaks were preceded by several years of increasing trap catches. However, the intensity of the outbreaks was not predicted by trapping alone. Trap catches preceding defoliation in 1986 were similar to trap captures prior to the 2000 outbreak; yet the intensity of the two outbreaks was very different. The outbreak in 1986 caused detectable

defoliation for one year, while defoliation in the 2000 outbreak was evident for three years. This confirms the need for additional population sampling, such as egg mass and larval sampling to help determine the intensity of outbreaks (Mason and Torgersen 1983, Kegley et al. 2004).

Cocoon and egg mass surveys are conducted in the fall of the same year that trap catches reach the threshold of 25 moths per trap. Larval surveys are performed in the spring and summer of the following year. All surveys are conducted in the vicinity of established plots or in other areas of concern. Cocoon and larval surveys provide estimates of population densities and give more accurate indications of outbreak potential and population trends. Pheromone trapping is designed to detect population changes over large geographic areas.

The DFTM, EWS **is not designed nor intended** to predict exactly where the defoliation will occur. Areas sampled on the ground are selected on the basis of the impact of potential DFTM defoliation on management objectives. Douglas-fir tussock moth EWS traps are **not** calibrated for use during an actual DFTM outbreak. As populations increase, a decline in trap catches will typically be noted. Once the traps have signaled a population increase, larval and cocoon/ egg mass surveys are used to determine population levels in that particular area (Sheehan and Ragenovich 2002).

Literature Cited

Daterman, G.E., R.L. Livingston, J.M. Wenz, and L.L. Sower. 1979. How to Use Pheromone Traps to Determine Outbreak Potential. USDA Agriculture Handbook No. 546.11 p.

Kegley, S.J., D. Beckman, and D.S. Wulff. 2004. 2003 North Idaho Douglas-fir Tussock Moth Trapping System Report. USDA Forest Service, Northern Region, Forest Health Protection Rpt. 04-6. 7pp.

Mason, R.R. 1978. Detecting Suboutbreak Populations Of The Douglas-fir Tussock Moth By Sequential Sampling Of Early Larvae In The Lower Tree Crown. General Technical Report PNW-238. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station. 9pp.

Mason, R.R. and T.R. Torgersen. 1983. Douglas-fir tussock moth handbook. How to Predict Population Trends. USDA Agriculture Handbook No. 610. 7 p.

Randall, C. 2002. Douglas-fir Tussock Moth Biological Evaluation, Palouse Ranger District, Clearwater National Forest, 2001. USDA Forest Service, Northern Region, Forest Health Protection Rpt. 02-5. 33pp.

Sheehan, K.A. and I. Ragenovich. 2002. Douglas-fir tussock moth early warning system trapping summary for Oregon and Washington. 2003. USDA Forest Service, Pacific Northwest Region, Forest Health Protection & Air Management Group/Natural Resources. 8p.




Figure 1. Map of plots trapped by IDL for DFTM in 2007 in northern Idaho.




Figure 2. Map of plots trapped by USFS for DFTM in 2007 in northern Idaho.




Figure 3. Mean trap catches of DFTM by IDL for plots north of Moscow, ID, from 1977 through 2007.




Figure 4. Aerially detected defoliation for the last two DFTM outbreaks in northern Idaho.

Appendix 1. Mean trap catch for IDL monitored plots from Coeur d'Alene to Moscow for the past 7 years.

IDL 2001 - 2007 Douglas-fir Tussock Moth Trap Results

| Plot # | Area | Mean Number of Moths Per Trap | | | | | | |
|--------|---------------------------------|-------------------------------|------|------|------|------|------|-------|
| | | 2007 | 2006 | 2005 | 2004 | 2003 | 2002 | 2001 |
| 3 | Lolo Pass | 0 ⁺ | 0 | 0 | 0 | 0 | 8.2 | 110.2 |
| 4 | Charles Butte | 0 ⁺ | 0 | 0 | 0 | 0.2 | 28.2 | 84.8 |
| 5 | Peterson Point | 0 ⁺ | * | 0 | 0 | 0.2 | 15.8 | 101.0 |
| 6 | East Dennis | 0.2 ⁺ | 0 | 0 | 0 | 1.2 | 75 | 101.2 |
| 7 | East Gold Hill | 0 ⁺ | 0 | 0 | 0 | 0.2 | 14.8 | 53.8 |
| 8 | Flat Creek | 0 ⁺ | 0.4 | 0 | 0.2 | 0 | 7.6 | 88.0 |
| 9 | Long Creek | 3 ⁺ | 0.2 | 0 | 0.2 | 0.2 | 33.6 | 0.2 |
| 10 | Paradise Point | 0.2 ⁺ | 0.2 | 0 | 0.2 | 0 | 17 | 91.8 |
| 11 | Mineral Mountain | 0.5 ⁺ | 0 | 0 | 0 | 1.8 | 75.2 | 56.4 |
| 12 | Mission Mountain | 0.2 ⁺ | 1.2 | 0 | 1.2 | 0.2 | 25.6 | 1.6 |
| 13 | Spring Valley Creek | 0 ⁺ | * | 0 | 0 | 0 | 5.4 | 58.0 |
| 14 | Vassar Meadows | 0.4 ⁺ | 0 | 0 | 0 | 0 | 95.8 | 102.8 |
| 15 | Fairview Knob | 0.4 ⁺ | 0 | 0 | 0 | 0.2 | 39 | 105.8 |
| 21 | West Twin (10-115) | 0.4 | * | 0 | 0 | 0 | 8.8 | 75.4 |
| 22 | Moscow Mtn (115-114) | 0 | 0 | 0 | 0 | 0.2 | 5.8 | 78.0 |
| 101 | Benewah | 0.2 ⁺ | 1.4 | 0 | 1.4 | 2.8 | 52.2 | 92.4 |
| 102 | Windfall Pass | 0.75 ⁺ | 0.6 | 0 | 0.6 | 0.6 | 40.4 | 99.6 |
| 103 | Squaw Creek | 0 | * | 0 | 0 | 0.2 | 9.4 | 89.2 |
| 104 | Moses Mountain | 0 | 0 | 0 | 0 | 0.2 | 6.4 | 67.8 |
| 105 | Little John Creek | 0.6 | 0 | 0 | 0 | 1.4 | 45 | 78.4 |
| 106 | Emida Peak | 0.4 | 0 | 0 | 0.2 | 2.6 | 64.2 | 75.8 |
| 107 | North-South Ski Area | 0 | 0 | 0 | 0 | 0.6 | 83.2 | 107.2 |
| 108 | Bald Mountain | * | * | 0 | 0 | 0 | 25.2 | 53.8 |
| 109 | Laird Park | 0 | 0 | 0 | 0 | 1 | 66 | 86.0 |
| 110 | North Fork Palouse River | 0 | 0 | 0 | 0 | 1 | 83.2 | 75.2 |
| 111 | Mica Mountain | 0.2 | 0 | 0 | 0 | 0.2 | 67.6 | 93.6 |
| 112 | Schwartz Creek | 0 | 0 | 0 | 0 | 0.2 | 80.6 | 110.6 |
| 113 | Big Bear Creek | 0.6 ⁺ | 0.6 | 0 | 0.6 | 0.2 | 47.8 | 87.0 |
| 114 | Big Meadow Creek | 0 ⁺ | 0.2 | 0 | 0.2 | 0 | 11.2 | 70.2 |
| 115 | East Twin Mountain | 0.4 ⁺ | 0.2 | 0 | 0.2 | 0 | 7.6 | 85.4 |
| 116 | Crane Point | 0 | * | 0 | 0 | 0 | 51 | 89.0 |
| 117 | Sheep Creek | 0 ⁺ | 0.2 | 0 | 0.2 | 0 | 27.8 | 83.2 |
| 118 | West Fork Mission Creek | 0.2 | * | 0 | 0 | 0 | 22.2 | 47.6 |
| 119 | 1 Mi N. of Mineral Mtn (11-216) | 0 | * | 0 | 0 | 0 | 25.2 | 0.2 |
| 200 | 2 mi W of Plummer | 2.6 | * | 0 | 0 | 0 | 16.2 | 80.2 |
| 201 | Coon Creek | 3 ⁺ | 2 | 0 | 0.4 | 0.2 | 21.6 | 93.8 |
| 202 | 3 mi E of Benewah | * ⁺ | 0.2 | 0 | 0.2 | 0.6 | 21 | 102.2 |
| 203 | Benewah Point | 0.4 | * | 0 | 0 | 0 | 8.2 | 92.4 |
| 204 | John's Point | * | * | 0 | 0 | 0 | 23.8 | * |
| 205 | 3 mi E of Charles Butte | 0.8 ⁺ | 0 | 0.2 | 0.2 | 0.4 | 63.6 | 72.6 |
| 206 | Sunset Mountain | * | * | 0 | 0 | 0 | 20.8 | * |
| 207 | West Fork Emerald Creek | 0 | * | 0 | 0 | 0 | 23.2 | * |
| 208 | Cedar Butte | 0 | * | 0 | 0 | 0 | 22.4 | 76.2 |
| 209 | Abes Knob | 0.2 | * | 0 | 0 | 0 | 23.8 | 88.4 |
| 210 | West Fork Deep Creek | 0.2 ⁺ | 0.2 | 0 | 0.2 | 0.2 | 77 | 90.6 |
| 211 | Cherry Butte | 0 ⁺ | 0 | 0 | 0.2 | 0.4 | 67.2 | 88.6 |
| 212 | Jackson Mountain | 0.2 | * | 0 | 0 | 0 | 19.6 | * |

IDL 2001 - 2007 Douglas-fir Tussock Moth Trap Results Cont.

| Plot # | Area | Mean Number of Moths Per Trap | | | | | | |
|--------|---|-------------------------------|------|------|------|------|-------|-------|
| | | 2007 | 2006 | 2005 | 2004 | 2003 | 2002 | 2001 |
| 216 | 1 mi NW of Mineral Mtn | 0 ⁺ | 0.4 | 0 | 0.4 | 0.2 | 1 | 0.2 |
| 217 | Head of Sheep Creek (216-117-2) | 0 ⁺ | 0.2 | 0 | 0.2 | 0.6 | 21.2 | 97.2 |
| 300 | Mission Mountain (#2) | 0 | 0.4 | 0 | 0.4 | 0.6 | 6.4 | 67.0 |
| 301 | 1.5 mi S of Mineral Mtn | 0 ⁺ | 0.2 | 0 | 0.2 | 0.2 | 69.4 | 91.2 |
| 302 | Middle Fork of Deep Creek 1 (301-216-1) | 1 | * | 0 | 0 | 0 | 63.8 | 3.6 |
| 303 | Middle Fork of Deep Creek 2 (301-216-3) | 0.2 | 0.4 | 0 | 0.2 | 1 | 58 | 15.8 |
| 400 | 3 mi S of Mineral Mtn | 0.6 ⁺ | 0.2 | 0 | 0.2 | 0.6 | 75.8 | 86.6 |
| 401 | Flynn Butte | 0 | 0 | 0 | 0 | 3.2 | 95.2 | 96.4 |
| 402 | 2 mi SE of Browns Mdw | 0.2 ⁺ | 0.2 | 0 | 0.2 | 0 | 15.2 | 57.4 |
| 500 | 3 mi SW of Harvard | 0 ⁺ | 0.2 | 0 | 0.2 | 0 | 18.8 | 74.6 |
| 501 | 3 mi S of Moon Hill | 0 | * | 0 | 0 | 0 | 16.2 | 97.6 |
| 502 | 3 mi W of Crane Point | 0.2 | * | 0 | 0 | 0.6 | 67.6 | 75.0 |
| 503 | 3 mi N of Stanford Point | 1 | * | 0 | 0 | 0 | 10.2 | 89.4 |
| 504 | 2 mi N of Stanford Point | 0 ⁺ | 0.4 | 0 | 0.4 | 0.2 | 47.8 | 86.2 |
| 505 | 1 mi SW of Stanford Point | 0.2 ⁺ | * | 0 | 0 | 0 | 38.4 | 47.0 |
| 506 | 1 mi S of Stanford Point | 1 | * | 0 | 0 | 0 | 23.4 | 67.8 |
| 507 | 1 mi NE of Stanford Point | 0 | 0 | 0 | 0 | 0.8 | 40.6 | 87.4 |
| 508 | 1 mi W of Stanford Point | 0.4 | 0.2 | 0 | 0.2 | 0 | 20.6 | 92.4 |
| 509 | 2 mi NW of Stanford Point | 1.2 ⁺ | 0.6 | 0.2 | 0.4 | 0.4 | 43.2 | 81.6 |
| 510 | Moon Hill | 0 ⁺ | 0.2 | 0 | 0.2 | 0.8 | 35 | 67.2 |
| 511 | 2 mi SE of Moon Hill | 0 | * | 0 | 0 | 0.2 | 13.2 | 80.4 |
| 512 | 3 mi S of Mineral Mtn | 0 | * | 0 | 0 | 0.2 | 70.2 | * |
| 513 | 2 mi SW of Moon Hill | 1.4 | * | 0 | 0 | 0 | 9.6 | 9.2 |
| 514 | 1.5 mi NW of Avon | 0 | * | 0 | 0 | 0 | 6.8 | 61.4 |
| 600 | 3.4 mi NNW of Princeton | 0.25 ⁺ | * | * | * | * | * | * |
| 601 | Macumber Meadows | 0 ⁺ | * | * | * | * | * | * |
| 602 | S of Shay Hill | 0.2 | * | * | * | * | * | * |
| 603 | 3 mi. S of Chatcolet | 0 | * | * | * | * | * | * |
| 701 | Fourmile Creek | 0.4 | * | 0 | 0 | 0 | 9 | 88.6 |
| 702 | North of Granite Point | 0 | * | 0 | 0.2 | 0 | 5.8 | 76 |
| 703 | Bergs Creek | 0 | * | 0 | 0 | 0 | 12.2 | 96.6 |
| 704 | West Fork Big Bear Creek | 0 ⁺ | 0.2 | 0 | 0.2 | 0.2 | 13.2 | 61 |
| 705 | 2 Mi NW of Stanford PT | 1.5 ⁺ | 0.8 | 0 | 0.8 | 0.4 | 46.4 | 89.4 |
| 706 | 1 Mi S. of Iron Mtn | 0.8 ⁺ | * | 0 | 0 | 0 | 27.2 | 87.8 |
| 707 | Iron Mtn | * | * | 0 | 0 | 0 | 6.6 | 97 |
| 708 | Little Bear Creek | 0.4 ⁺ | * | 0 | 0 | 0 | 65.6 | 108.6 |
| 709 | Ruby Creek | 0 | * | 0 | 0 | 0 | 50.4 | 96.2 |
| 710 | Turnbow Creek | 2.4 ⁺ | 1.4 | 0 | 1.4 | 0.2 | 43 | 70.6 |
| 711 | East Fork Flat Creek | 2 ⁺ | 2.6 | 0 | 2.6 | 0.2 | 55 | 71.4 |
| 712 | Turnbow Point | 0.2 | * | 0 | 0 | 0.2 | 7.8 | 38 |
| 713 | 3 Mi S. of Potlatch | 0 ⁺ | * | 0 | 0 | 0 | 6.6 | 30 |
| 714 | Rocky Point | 0.8 | * | 0 | 0 | 0 | 13.2 | 79.6 |
| 715 | Hatter Creek | 0 ⁺ | 0.6 | 0 | 0.6 | 0.2 | 7.4 | 32 |
| 716 | Head of Hatter Creek | 0 | * | 0 | 0 | 0 | 11.8 | 80.8 |
| 717 | Nora Creek | 0 | * | 0 | 0 | 0 | 21.2 | 81.4 |
| 718 | Crummaring Creek | 0.2 | * | 0 | 0 | 0 | 12.4 | 70.4 |
| 719 | Basalt Hill | 0.2 | * | 0 | 0 | 0 | 19 | 11.6 |
| 720 | Browns Meadow | 0.4 | 0 | 0 | 0 | 0.2 | 11.2 | 2.6 |
| 721 | Smith Creek | 0 | * | 0 | 0 | 0 | 100.2 | 70.6 |

IDL 2001 - 2007 Douglas-fir Tussock Moth Trap Results Cont.

| Plot # | Area | | Mean Number of Moths Per Trap | | | | | | |
|-----------------------------------|-----------------------------|----------------|-------------------------------|------|------|------|-------|-------|------|
| | | | 2007 | 2006 | 2005 | 2004 | 2003 | 2002 | 2001 |
| 722 | Prospect Peak | 0 | * | 0 | 0 | 0 | 31.2 | 56.8 | |
| 723 | West Fork Mission Creek | 0 | * | 0 | 0 | 0 | 27.8 | 22.2 | |
| 724 | Huckleberry Mtn | 0 [†] | * | 0 | 0 | 0 | 16.6 | 77.2 | |
| 725 | North Fork Pine Creek | 0.75 | * | 0 | 0 | 0 | 21.6 | 93 | |
| 726 | Mineral Creek | 0 | * | 0 | 0 | 0 | 20.2 | 78 | |
| 727 | South of Sanders | 0 | * | 0 | 0 | 0 | 77.8 | 86.8 | |
| 800 | Mason Butte | 7.25 | * | * | * | 0 | 20.8 | 63 | |
| 801 | 1 mi SW of Moctileme Butte | 0.2 | * | * | * | 0 | 30.2 | 91.4 | |
| 802 | 1.9 mi S of Plummer | 0 | * | * | * | 0 | 24.8 | 75.2 | |
| 803 | Little Plummer Creek | 5.8 | * | * | * | 0 | 18 | 54.4 | |
| 804 | Syringa Creek | 0 | * | * | * | 0 | 21.2 | 66.4 | |
| 805 | John Point | * | * | * | * | 0 | 20.4 | 61.6 | |
| 806 | 2 mi W of Pettis Point | 0 | * | * | * | 0 | 22.6 | 71.2 | |
| 807 | Davis Creek | 0 | * | * | * | 0 | 17.8 | 55.6 | |
| 808 | Renfro Creek | 0 | * | * | * | 0 | 14.8 | 44.2 | |
| 809 | Crystal Creek | 0.2 | * | * | * | 0 | 10.4 | 29.4 | |
| 810 | Child Creek | 0 | * | * | * | 0 | 17.2 | 52.8 | |
| 811 | Hobo Pass | 0.6 | * | * | * | 0 | 7.8 | 25.4 | |
| 812 | Hemlock Butte | 0.4 | * | * | * | 0 | 9.2 | 28.2 | |
| 813 | Carpenter Peak | 0 | * | * | * | 0 | 18.8 | 57.8 | |
| 814 | Tyson Creek | 0 | * | * | * | 0 | 30.2 | 87.6 | |
| 815 | Heinaman Creek | 0 | * | * | * | 0 | 25.2 | 85.2 | |
| 816 | Green Mtn | 0 | * | * | * | 0 | 31 | 86.2 | |
| 817 | Willow Creek | 1.2 | * | * | * | 0 | 22.2 | 73.2 | |
| 818 | Head of Emerald Creek | 0.6 | * | * | * | 0 | 28.2 | 86 | |
| 819 | East Fork Emerald Creek | 0 | * | * | * | 0 | 25 | 75.2 | |
| 820 | Head of Bobs Creek | 0 | * | * | * | 0 | 25.4 | 79 | |
| 821 | East Fork of Potlatch River | 0 | * | * | * | 0 | 25.2 | 67.2 | |
| 822 | Head of Moose Creek | 0.2 | * | * | * | 0 | 24.8 | 69.6 | |
| 823 | Beals Butte | 0 | * | * | * | 0 | 39 | 106.2 | |
| Number of Sites Trapped: | | 120 | 51 | 98 | 98 | 122 | 122 | 117 | |
| Average Number of Moths per Plot: | | 0.42 | 0.33 | 0.00 | 0.16 | 0.23 | 31.30 | 71.50 | |

* Indicates Sites Not Trapped

[†] Indicates larval survey

Appendix 2. Mean trap catch for USFS monitored plots from Potlatch to Lucille for the past 7 years.

USFS 2001 - 2007 Douglas-fir Tussock Moth Trap Results

| Plot # | Site Name | Mean Number of Moths per Trap | | | | | | |
|--------------------------------|---------------------|-------------------------------|------|------|------|------|------|------|
| | | 2007 | 2006 | 2005 | 2004 | 2003 | 2002 | 2001 |
| 1-1 | Lodge Pt | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 1.2 | 1.6 |
| 1-2 | Goddard | * | * | * | * | | * | * |
| 1-3 | Pine Knob | 0.2 | 0.3 | 0.0 | 0.0 | 0.0 | 1.0 | 4.8 |
| 1-4 | Potatoe Hill | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.2 |
| 1-5 | Big Tinker | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.6 | 1.4 |
| 2-1 | Rhett Cr | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 |
| 2-2 | Christie Cr | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 8.0 |
| 2-3 | Cow Cr Saddle | * | * | 0.0 | 0.0 | 0.0 | 0.2 | 0.2 |
| 2-4 | Low Saddle | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 |
| 2-5 | S. Cow Cr | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.8 |
| 2-6 | Spring Mtns | 0.0 | 0.0 | * | * | * | * | * |
| 3-1 | Keuterville | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.2 |
| 3-2 | Cottonwood Butte | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.8 |
| 4-1 | Lake Waha | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 10.2 |
| 4-2 | Black Pine | 0.2 | 0.0 | 0.0 | 0.0 | | 0.2 | 18.2 |
| 4-3 | Junction | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | * |
| 4-4 | Captain John | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 3.6 |
| 4-5 | Webb Cr | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.4 |
| 4-6 | Forest | * | * | * | * | * | * | * |
| 5-1 | Johnson | * | 0.0 | 0.0 | 0.0 | 0.0 | 4.8 | 4.0 |
| 5-2 | Angel Butte | * | 0.0 | 0.0 | 0.0 | 0.4 | 0.8 | 5.8 |
| 5-3 | Grangemont | 1.4 | 1.4 | 0.0 | 0.0 | 0.4 | 2.2 | 16.2 |
| 5-4 | Bargamin Cr | 4.6 | 0.0 | 0.0 | 0.0 | 0.0 | 4.8 | 35.6 |
| 5-5 | Bald Mtn | 3.4 | 1.8 | 0.0 | 0.0 | 0.2 | 9.0 | 36.0 |
| 5-6 | Summit Landing | 3.2 | 0.6 | 0.0 | 0.0 | 0.2 | 0.0 | 14.6 |
| 5-7 | Shin Pt | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 13.2 |
| 5-8 | Swanson Cr | 0.8 | 0.6 | 0.0 | 0.0 | 1.4 | 0.0 | 17.5 |
| 5-9 | Skull Cr | * | * | * | * | | * | * |
| 5-10 | Cooper | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 3.8 |
| 6-1 | Canyon Junction | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 11.2 |
| 6-2 | Fan saddle | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.6 |
| 7-1 | Laird Park | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 52.2 | * |
| 7-2 | Little Bald Mtn. | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 22.0 | * |
| 7-3 | Little Boulder Cr. | 0.0 | 1.2 | 0.0 | 0.0 | 4.0 | 40.4 | * |
| 7-4 | W. Fork Potlatch R. | 0.0 | 0.8 | 0.6 | 0.0 | 2.4 | 40.4 | * |
| 7-5 | Elk Creek Falls | 0.2 | 0.4 | 0.4 | 0.0 | 4.8 | 15.8 | * |
| 7-6 | Morris Creek | 0.0 | 0.2 | 0.0 | 0.0 | 0.2 | 26.5 | * |
| Number of Plots Trapped: | | 31 | 33 | 33 | 33 | 32 | 33 | 26 |
| Mean Number of Moths per Site: | | 0.47 | 0.24 | 0.03 | 0.01 | 0.45 | 6.82 | 8.30 |
| * Indicates Plots Not Trapped | | | | | | | | |